

Vector Symmetries and Θ -Vacuum

Vicente Azcoiti and Angelo Galante

*Departamento de Física Teórica, Facultad de Ciencias, Universidad de Zaragoza,
50009 Zaragoza (Spain).*

ABSTRACT

Under very general assumptions we show that Vafa-Witten theorem on vector symmetries in vector-like theories can be extended to some physically relevant gauge theories with non-positive definite integration measure as QCD with a Θ -vacuum term.

The vacuum realization of vector and axial symmetries in gauge theories with dynamical fermions, as QCD , plays a fundamental role in the understanding of the dynamics of these models at zero and finite temperature. The spontaneous breakdown of chiral symmetry in QCD at zero temperature and the realization of the Goldstone theorem allow us to understand many essential features of the low energy spectrum of this model, and the restoration of this symmetry at high temperature is the main signal expected for the deconfining phase transition.

Concerning vector symmetries as flavor or baryon number, Vafa and Witten showed [1] few years ago that they can not be spontaneously broken in vector-like theories as QCD . The two main ingredients in the demonstration were *i*) the positivity of the path integral measure in the effective gauge theory obtained after integration of the Grassman degrees of freedom and *ii*) the antihermiticity and local character of the massless Dirac operator.

The second ingredient was essential in order to derive bounds which show exponential fall-off at large distances for the massive quark propagator in any background gauge field. The authors of [1] noticed that the correlation function of any quark-composite local gauge-invariant operator carrying a non vanishing conserved vector charge should be proportional to at least one power of the quark propagator. Taking also into account that the mean value of an upper-bounded quantity computed with a probability distribution function is always less or equal than its upper-bound, an exponential fall-off at large distances was found for the correlation function of all vector charge carrying operators. This result allowed to show not only that the mean

value of any local order parameter for a vector symmetry should vanish but also that massless quark composites can not be constructed from massive constituents.

Even if Vafa-Witten theorem applies also to vector-like theories at finite temperature, there are some physically relevant formulations in which at least one of the two essential ingredients in the demonstration are not realized. The most interesting cases are QCD with a Θ -vacuum term and QCD at finite chemical potential.

In QCD with a Θ -vacuum term the factor of i picked-up by the $F\tilde{F}$ operator under Wick rotation makes the euclidean action complex. Even if the partition function is real, the integration measure is no longer positive definite and therefore the first ingredient in the theorem breaks-down. The second one is however preserved since the Θ -vacuum term does not change the structure of the massless Dirac operator.

The introduction of a finite chemical potential in $SU(N)$ gauge theories makes also the effective gauge action obtained after integration of the Grassman degrees of freedom complex in the most interesting case of $N = 3$. Furthermore the introduction of a chemical potential modifies also the structure of the massless Dirac operator. It loses its antihermiticity and the second ingredient of the theorem breaks-down too.

Since both cases, QCD with a Θ -vacuum term and QCD at finite chemical potential, are physically relevant, it is worthwhile to ask to which extent and under what limitations the theorem can be extended to these formulations. We will devote this paper to QCD with a Θ -vacuum term leaving

the finite chemical potential case, which shows the additional complication previously stated, for a separate publication.

In order to extend the theorem to QCD with a Θ -vacuum term we will made use of two very general assumptions. First we will assume that Parity is not spontaneously broken in QCD at $\theta = 0$, which is a necessary condition for the euclidean free energy density be well defined at $\theta \neq 0$ [2].

The second general assumption, which is standard in statistical mechanics, is that in the thermodynamical limit any intensive operator does not fluctuate in a vacuum or equilibrium state. This is equivalent to the general Quantum Field Theory statement that all connected correlation functions of local operators verify cluster.

To better understand the physical meaning of this last assumption let us think for a while in the Ising model. In the high temperature symmetric phase the system has one vacuum or equilibrium state. The density of magnetization is an intensive operator and its probability distribution function will become a single δ function in the thermodynamical limit since otherwise we would have fluctuations of the density of magnetization. Since the $Z(2)$ symmetry of the Ising model is realized in the high temperature phase, the δ function is centered at the origin.

In the low temperature phase the $Z(2)$ symmetry is spontaneously broken. Since the density of magnetization m_I is an order parameter for this symmetry, its probability distribution function becomes now $1/2$ the sum of two δ functions. The density of magnetization fluctuates in the thermodynamical limit because we have two vacuum or equilibrium states and m_I is

not invariant under $Z(2)$ transformations. However if we choose an intensive operator invariant under $Z(2)$ transformations, as for instance the energy density or m_I^2 , it does not fluctuate in spite of the fact that we have two vacuum states. Indeed the mean value of any power of these operators does not depend on the vacuum state it is computed.

The lesson we learn from this simple example is that if the only reason to have a degenerate vacuum is spontaneous symmetry breaking (*i.e.* no accidental vacuum degeneration) and we choose an intensive operator invariant under symmetry transformations, it does not fluctuate in the thermodynamical limit independently of the symmetry realization. These features are not exclusive of the Ising model but a general property of any statistical system.

Let us consider now the euclidean formulation of QCD with a Θ -vacuum term. The euclidean partition function is

$$\mathcal{Z} = \int [dA_\mu][d\bar{\psi}][d\psi] e^{-\int d^4x (\mathcal{L}(x) - \frac{i\theta}{16\pi^2} X(x))} \quad (1)$$

where $\mathcal{L}(x)$ is the standard QCD lagrangian and $X(x) = \epsilon_{\mu\nu\alpha\beta} \text{Tr} F_{\mu\nu} F_{\alpha\beta}$ the Θ -vacuum lagrangian. $X(x)$ is real in euclidean space and we have exhibited in (1) the factor of i which arises from Wick rotation.

When we integrate out the fermionic degrees of freedom we get an effective gauge theory

$$\mathcal{Z} = \int [dA_\mu] \det \Delta(m, A_\mu) e^{-S_G(A_\mu)} e^{i\theta \hat{X}(A_\mu)}, \quad (2)$$

where $\Delta(m, A_\mu) = m + i\Lambda(A_\mu)$ is the Dirac operator for quarks of bare mass m , $S_G(A_\mu)$ the pure gauge action and $\hat{X}(A_\mu) = \frac{1}{16\pi^2} \int d^4x X(x)$ the Θ -vacuum action.

The integrand in (2) is a complex number due to the Θ -vacuum contribution. The imaginary part of the integrand gives no contribution to the partition function (the QCD action is parity invariant). The real part however is not positive definite except at $\theta = 0$.

Let $J(x)$ be any quark composite gauge invariant operator with non zero isospin or baryon number and let us consider the correlation function $Q(x) = \langle J(x)J^*(0) \rangle$. This correlation function will be given by the following ratio

$$Q(x) = \frac{\int [dA_\mu] Q_A(x) \det \Delta(m, A_\mu) e^{-S_G(A_\mu) + i\theta \hat{X}(A_\mu)}}{\int [dA_\mu] \det \Delta(m, A_\mu) e^{-S_G(A_\mu) + i\theta \hat{X}(A_\mu)}} \quad (3)$$

where $Q_A(x)$ is the correlation function in a given background gauge field $A_\mu(x)$. If, as previously assumed, $J(x)$ carries a conserved fermion charge, then $Q_A(x)$ will be proportional to at least one power of the fermion propagator $\Delta_A^{-1}(m, 0, x)$ from 0 to x . Vafa and Witten essentially showed in [1] that the norm of the quark propagator matrix

$$\|\Delta_A^{-1}(m, 0, x)\| \leq \alpha e^{-\beta \|x-y\|} \quad (4)$$

falls-off at least exponentially for any gauge configuration since α and β in (4) are two constants independent of the gauge configuration. In the previous statement the word essentially stands for the fact that, strictly speaking, relation (4) is not true in the continuum formulation (it is true in the lattice regularization approach). This problem can be surmounted by considering smeared correlation functions [1] or using a lattice regularization scheme. In order to preserve as much as possible the simplicity in the notations, we will assume relation (4) true in what follows.

At $\theta = 0$, relation (4) plus the fact that the integration measure in (3) defines a probability distribution function, carried Vafa and Witten to conclude that a bound similar to (4) holds also for the correlation function $Q(x)$ and consequently that vector symmetries can not be spontaneously broken. They noticed also in [1] that introduction of a Θ -vacuum term would invalidate the proof because the positivity of the integration measure is loosed.

To overcome this problem, we will make use now of the second general assumption discussed in the introduction of this paper, *i.e.* in any statistical system intensive operators do not fluctuate or equivalently, in a well defined quantum system, all the correlation functions of local operators verify the cluster property. We will apply this general assumption to the intensive operator $\tilde{X}(A_\mu) = \frac{1}{V} \frac{1}{16\pi^2} \int d^4x \hat{X}(A_\mu)$ which defines the Θ -vacuum term in the action.

The cluster property tells us that

$$\langle \tilde{X}^n \rangle = \langle \tilde{X} \rangle^n \quad \forall n \quad (5)$$

Due to the factor of i which arises from Wick rotation, the mean value $\langle \tilde{X} \rangle$ is a pure imaginary number whereas the operator $\tilde{X}$ takes real values in euclidean space. This two features however are not in contradiction since the mean value in (5) is computed with a complex action. In order to avoid complications related to factors of i , we will consider the operator

$$Y = \tilde{X}^4, \quad (6)$$

the mean value of which is always real and positive. Furthermore (5) implies

analogous relations for the Y operator:

$$\langle Y^n \rangle = \langle Y \rangle^n \quad \forall n \quad (7)$$

Let us assume now that O is any parity conserving operator as for instance the operator Y previously defined or the correlation function of any gauge invariant quark-composite operator driving flavor or baryon number. We can write

$$\langle O \rangle = \frac{\int [dA_\mu] [d\bar{\psi}] [d\psi] O(\bar{\psi}, \psi, A_\mu) e^{-S_{QCD}(\bar{\psi}, \psi, A_\mu)} \cos(\theta V \tilde{X})}{\int [dA_\mu] [d\bar{\psi}] [d\psi] e^{-S_{QCD}(\bar{\psi}, \psi, A_\mu)} \cos(\theta V \tilde{X})} \quad (8)$$

which, after integration of the Grassman degrees of freedom becomes

$$\langle O \rangle = \frac{\int [dA_\mu] O(A_\mu) \det \Delta(A_\mu) e^{-S_G(A_\mu)} \cos(\theta V \tilde{X})}{\int [dA_\mu] \det \Delta(A_\mu) e^{-S_G(A_\mu)} \cos(\theta V \tilde{X})} \quad (9)$$

where, in order to leave notation as simple as possible, we have used the same symbol O to design the operator in the effective gauge theory obtained after integration of the fermion degrees of freedom.

In order to define a pseudo-probability distribution function we will introduce in (9) the trivial identity

$$1 = \int_0^\infty d\bar{Y} \delta(Y(A_\mu) - \bar{Y}) \quad (10)$$

which allows us to write expression (9) as

$$\langle O \rangle = \frac{\int_0^\infty d\bar{Y} N(\bar{Y}) O(\bar{Y}) e^{-S_{eff}(\bar{Y})} \cos(\theta V \bar{Y}^{1/4})}{\int_0^\infty d\bar{Y} N(\bar{Y}) e^{-S_{eff}(\bar{Y})} \cos(\theta V \bar{Y}^{1/4})} \quad (11)$$

where $N(\bar{Y})$ is the density of states

$$N(\bar{Y}) = \int [dA_\mu] \delta(Y(A_\mu) - \bar{Y}) \quad (12)$$

and $e^{-S_{eff}(\bar{Y})}$ and $O(\bar{Y})$ are respectively

$$e^{-S_{eff}(\bar{Y})} = \frac{\int [dA_\mu] \det \Delta(A_\mu) e^{-S_G(A_\mu)} \delta(Y(A_\mu) - \bar{Y})}{\int [dA_\mu] \delta(Y(A_\mu) - \bar{Y})}$$

$$O(\bar{Y}) = \frac{\int [dA_\mu] \det \Delta(A_\mu) e^{-S_G(A_\mu)} O(A_\mu) \delta(Y(A_\mu) - \bar{Y})}{\int [dA_\mu] \det \Delta(A_\mu) e^{-S_G(A_\mu)} \delta(Y(A_\mu) - \bar{Y})} \quad (13)$$

This relations allow us to write for $\langle O \rangle$ the simple expression

$$\langle O \rangle = \int_0^\infty d\bar{Y} P(\bar{Y}) O(\bar{Y}) \quad (14)$$

where

$$P(\bar{Y}) = \frac{N(\bar{Y}) e^{S_{eff}(\bar{Y})} \cos(\theta V \bar{Y}^{1/4})}{\int_0^\infty d\bar{Y} N(\bar{Y}) e^{S_{eff}(\bar{Y})} \cos(\theta V \bar{Y}^{1/4})} \quad (15)$$

is a pseudo-probability distribution function since it is not positive definite due to the factor $\cos(\theta V \bar{Y}^{1/4})$ in the numerator. In spite of that, equations (7) and (14) tell us that in the infinite volume limit

$$\int_0^\infty d\bar{Y} P(\bar{Y}) \bar{Y}^n = \bar{Y}_0^n \quad \bar{Y}_0^n \geq 0 \quad (16)$$

Equation (16) implies that in the infinite volume limit $P(\bar{Y})$ becomes a true probability distribution function: a Dirac distribution $\delta(\bar{Y} - \bar{Y}_0)$. The mean value of operator O will be in this limit

$$\langle O \rangle = \int_0^\infty d\bar{Y} \delta(\bar{Y} - \bar{Y}_0) O(\bar{Y}) = O(\bar{Y}_0) \quad (17)$$

where $\bar{Y}_0$ will depend on the theory parameters: gauge coupling, quark masses and θ angle.

It turns out that to compute the expectation value of our parity invariant operator it is enough to calculate its average over a subset of gauge fields, actually all the gauge fields characterized by a fixed value of $Y(A_\mu)$. The

expectation value of our operator coincides with its mean value given by the expression (13) and evaluated at $\bar{Y} = \bar{Y}_0$.

Let's see the consequences for any correlation function of a gauge invariant composite quark operator driving flavor or baryon number. It is clear that expression (13) defines a true probability distribution function (the integration measure is positive definite for every $\bar{Y}$). It follows that if $|O(A_\mu)|$ verifies the Vafa-Witten bounds for all gauge configurations the same happens for $|O(\bar{Y})|$ whatever the value of $\bar{Y}$. In particular this will be true for $\bar{Y} = \bar{Y}_0$ and therefore we conclude that the expectation value of any of these correlation functions in presence of a non zero Θ -vacuum term shows exponential fall-off at large distances.

In conclusion the two main consequences of Vafa-Witten theorem *i)* the impossibility to break spontaneously a vector symmetry and *ii)* the non existence of massless bound states made up with massive constituents, apply also to QCD with a Θ -vacuum term, assumed the euclidean formulation of this model can be consistently done.

Acknowledgements

This work has been partially supported by CICYT (Proyecto AEN97-1680). A.G. was supported by a Istituto Nazionale di Fisica Nucleare fellowship at the University of Zaragoza.

References

- [1] C. Vafa, E. Witten, Nucl. Phys. **B234** (1984) 173.
- [2] V. Azcoiti, A. Galante, hep-th/9901068.